

# Mars Seismic Catalogue, InSight Mission; V8 2021-10-01

(actual release date: 2021-10-06)

Authors: John Clinton, Fabian Euchner and the MQS frontline team

## Overview

This is the description of version 8 (V8) of the Marsquake Catalogue for InSight, which includes the Martian seismic events up to June 30, 2021 / Sol 922 as identified by the InSight Marsquake Service (MQS). The catalogue files are available at IPGP and IRIS. New versions of the catalogue are released in sync with updated waveform data releases.

The citation for the catalogue is:

**InSight Marsquake Service (2021). Mars Seismic Catalogue, InSight Mission; V8 2021-10-01. ETHZ, IPGP, JPL, ICL, Univ. Bristol. <https://doi.org/10.12686/a13>**

This catalogue is an update of V1-7 (*InSight Marsquake Service, 2020a, b, c, d; 2021a, b, c*).

The catalogue is provided in two files, both in QuakeML format. One is in standard QuakeML 1.2 format, known as “basic event description” (BED). Thus, it validates against the QuakeML 1.2 schema. The second includes two Mars-specific extensions in separate XML namespaces. These include basic information for single station locations and Mars catalogue management, which is not available in the standard BED format. The additional information includes: distance, back azimuth, Mars event type, Mars event quality, and Marsquake name. The XML schema of the Mars-specific extension is provided in V4 and is unchanged since then.

A detailed description of the V3 version of the catalogue, as well as key event presentations and MQS procedures, is provided at *Clinton et al. (2021)*. All MQS conventions for this version, V8, are unchanged from this publication. Key details are also repeated here.

The software that has been used to compile this version (V8) is the same as released in V5. The code of the basic GUI used by MQS, developed by ETH and gempa GmbH, is available at zenodo.org ([doi:10.5281/zenodo.4033316](https://doi.org/10.5281/zenodo.4033316)). This software provides the framework for data review, Marsquake detection, identification and quantification. The code to compute MQS distances of events, given P and S picks, is available at zenodo.org ([doi:10.5281/zenodo.4302312](https://doi.org/10.5281/zenodo.4302312)). Updates and/or additional software critical for MQS operations, will continue to be provided in association with future catalogue releases when relevant.

An overview of the major changes between V7 and V8 (this version) is at the end of this document; detailed changelogs for each of the two catalogue files are provided separately.

## MQS conventions

MQS assigns an event type and quality to each seismic event. The event type reflects the frequency content. The event quality is assigned based on the signal strength and ability to identify and interpret the phase arrivals.

### MQS Event Type

<b>Low Frequency family: event energy generally at long period</b>	
Low frequency (LF)	energy in 3 components all below 2.4Hz.
Broadband (BB)	energy in 3 components predominantly below 2.4Hz though also includes excitement at and possibly above 2.4Hz.
<b>High Frequency family: event energy generally at high frequency</b>	
High Frequency (HF)	energy in 3 components predominantly at 2.4Hz and above. ‘Predominantly’ indicates some energy below 2.4Hz is possible.
2.4Hz	energy in 3 components centered around 2.4Hz resonance, with very limited excitation above or below. (It is likely these are small amplitude HF events.)
Very High Frequency (VF)	special case of high frequency events that show clear differences in energy between vertical and horizontal components. Horizontal energy is significantly larger than vertical energy at higher

	frequencies.
<b>Other Signals</b>	
Super High Frequency (SF)	very short duration high frequency events that do not include energy at 2.4Hz or below. Typically between 5-10Hz, and horizontal energy is significantly larger than vertical energy.

### MQS Event Quality

Label	Quality summary	Key features
A	High	Multiple clear and identifiable phases / clear polarisation (implies possibility both distance and back azimuth are determined, and hence location)
B	Medium	Multiple clear and identifiable phases but no polarisation (implies possibility of distance but no location) OR polarisation, but not enough clear phase picks for a distance estimate
C	Low	Signal is clearly observed but phase picking is challenging: <ul style="list-style-type: none"> <li>- (HF/2.4Hz/VF) Pg and Sg pickable, but speculative OR large uncertainty OR low SNR</li> <li>- (LF/BB) no clear phases can be identified OR only a single phase is clearly identifiable OR multiple phases are identifiable, but no clear picks can be attributed to P and S phases</li> <li>- (SF) peak signal amplitude of data with 7 - 9Hz BP filter above <math>2 \times 10^{-9}</math> m/s</li> </ul>
D	Suspicious	<ul style="list-style-type: none"> <li>- Signal only weakly observed OR</li> <li>- Signal may not be attributable to a seismic event OR</li> <li>- (HF/2.4/VF) impossible to pick both Pg and Sg OR</li> <li>- (SF) peak signal amplitude of data with 7.9Hz filter is below <math>2 \times 10^{-9}</math> m/s</li> </ul>

### MQS Event Names

Events belonging to the Low and High Frequency families are labelled following the convention S[xxxx][z]; where [xxxx] indicates the InSight mission sol the event begins on (starting from sol 0, the sol InSight landed on Mars), and [z] is a letter that ensures unique names if multiple events occur on a single Sol.

SF events are assigned the prefix letter T instead of S in order to clearly separate them from other events: T[xxxx][z].

### MQS Phase Picks

Onset Phase Picks: When possible, MQS selects the first arrival times for distinct energy packets. Pick time uncertainties are on the order of seconds, if made on the waveform in the time domain; and on the order of 10's of seconds, if these are based on a distinct new signal visible on a spectrogram. Typically, only 1 or 2 energy packets are identified, if any, and are labelled P and S for LF/BB event types, and Pg and Sg for HF, VF and 2.4Hz event types. In rare cases, when phases cannot be clearly attributed to P or S, they are labelled x1, x2, x3...

SF events do not have phase assignments.

For each event, MQS also includes 'picks' for event start and end and start and end of noise windows with similar noise as observed during the event. Since there are often numerous glitches occurring within the event time window, we also include 'clean', glitch-free P and S coda windows when possible. Depending on the event type, the time at which peak amplitudes occur with bandpassed signals are also indicated. MQS is tracking all significant glitches within the event start and end window, but these are currently not provided in the catalogue.

Pick uncertainties are assigned for P/S/Pg/Sg/x? but not for any other pick type.

### Distances, Back Azimuth and Location

**BB/LF events:** If multiple picks are assigned as P and S phases, a distance is estimated using a priori Martian velocity models. The back-azimuth can be estimated using the first phase arrival, assumed to be P, if polarization is present. A single station location estimate can be made by combining the distance and back-azimuths. This approach is based on *Böse et al. (2016)*, and outlined in *Clinton et al. (2021)*. Distance / back-azimuth / location uncertainties are included in the catalogue.

HF, VF and 2.4Hz events: If multiple picks are assigned as Pg and Sg phases a preliminary distance estimate is made using a simple crustal velocity model with  $V_s=2.3\text{km/s}$ ,  $V_p/V_s=1.73$ . There are no back-azimuth estimates for any of these events. Location uncertainty is provided as  $\pm(0.75 \times \text{Distance})$  ([van Driel et al., 2021](#)).

SF events: there are currently no distance or back-azimuth estimates for these events ([Dahmen et al., 2020](#)).

Only a handful of events in the catalogue include a computed latitude/longitude location. A location is required for a valid QuakeML origin, so by default all other events are assigned the location of the lander, at  $\text{lat}=4.5024^\circ$ ,  $\text{lon}=135.6234^\circ$ .

### Depth

Depths are not regularly included in the V8 catalogue (unchanged from previous catalogues). The only exceptions are two (S0235b, S0173a) of the three events for which moment tensors from Brinkman et al. (2021) are listed in the catalogue. The origins that have been used to compute the moment tensors are included and they also contain the best-fitting hypocentral depth (in metres). Note these are not the preferred origins. These depth values are new in V8.

### Magnitude

Catalogue version V8 uses new magnitude relations first introduced in V7, as described in [Böse et al. \(2021\)](#), replacing magnitudes in previous catalogues that were based on a pre-landing study. All events that have catalogue distances are assigned a  $M_w^{Ma}$ . Magnitude scales using P and S ( $m_b^{Ma}$  and  $m_{bS}^{Ma}$ ) body phase amplitudes, 2.4Hz resonance amplitudes ( $M_{2.4\text{Hz}}^{Ma}$ ), and spectral fitting ( $M_{Wspec}^{Ma}$ ) are included, when possible. Magnitude uncertainties are included.

For the Low Frequency family BB events that have both multiple origins based on S-P and 'aligned' (see below) distance estimates, magnitudes are provided for each distance estimate.

## Alternative Event Information

Other groups within the InSight Science team are contributing pick and location information that are being included in the MQS catalogue. These include:

### Alignments: Phase Picks, Locations, Origin Times

[Giardini et al. \(2020\)](#) introduce a procedure that provides aligned epicentral distances for good quality LF/BB events that is based on similarity of waveform envelopes. Metadata includes not just the epicentral distance but also an associated origin time as well as aligned P and S pick times. Information from this alignment procedure has been provided in the catalogue since V3. Since V7 these have been updated to include the new LF/BB events and adopting a new background velocity model from [Stähler et al. \(2021\)](#). Aligned distances have a methodID attribute of `smi:insight.mqs/algorithms/distance/aligned` in their corresponding DistanceComputation element, whereas S-P distances have a methodID attribute of `smi:insight.mqs/algorithms/distance/S-P_phases`.

Aligned distances are given as the preferred distance when they are available.

### Secondary Phase Picks

Various groups within the InSight Science teams but independent of MQS have proposed secondary body phase and core phase picks ([Stähler et al., 2021](#); [Khan et al., 2021](#)). Since different approaches are required, these also include different P and S picks. These have been added to the V7 catalogue as additional pick elements and are unchanged in V8. These picks are not associated to an origin through a corresponding arrival element.

### Moment Tensors

Since V7 moment tensor solutions as described in Brinkman et al. (2021) have been included. These are unchanged in V8 except for the addition of hypocentral depth for events S0235b and S0173a. The moment tensor information is contained in focal mechanism XML elements.

## V8 Catalogue Overview

Marsquake type events (Number in brackets is the increase since V7)

	<b>Total</b>	<b>A</b>	<b>B</b>	<b>C</b>	<b>D</b>
<b>Total</b>	671 (+158)	5 (+1)	111 (+15)	256 (+55)	299 (+87)
LF	42 (+5)	3	7 (+1)	17 (+3)	15 (+1)
BB	19 (+3)	2 (+1)	5 (+1)	10 (+1)	2
HF	63 (+6)	-	38 (+5)	19 (+1)	6
2.4Hz	501 (+135)	-	44 (+5)	188 (+45)	269 (+85)
VF	46 (+9)	-	17 (+3)	22 (+5)	7 (+1)

### Super high frequency events

	<b>Total</b>	<b>A</b>	<b>B</b>	<b>C</b>	<b>D</b>
SF	953 (+116)	-	-	232 (+54)	721 (+62)

### Overview of Major Changes from V7 to V8

None

## References

- Böse, M. et al., 2016.** A Probabilistic Framework for Single-Station Location of Seismicity on Earth and Mars. PEPI. doi.org/10.1016/j.pepi.2016.11.003
- Böse, M. et al., 2021.** Magnitude Scales for Marsquakes Calibrated from InSight Data, BSSA. doi.org/10.1785/0120210045
- Brinkman, N. et al., 2021.** First Focal Mechanisms of Marsquakes, JGR Planets. doi.org/10.1029/2020JE006546
- Clinton, J. et al., 2021.** The Marsquake Catalogue from InSight, Sols 0-478. doi.org/10.1016/j.pepi.2020.106595
- Dahmen, N. et al., 2020.** Super high frequency events: a new class of events recorded by the InSight seismometers on Mars. JGR Planets. doi.org/10.1029/2020JE006599
- Giardini, D. et al., 2020.** The seismicity of Mars, Nature Geoscience. doi:10.1038/s41561-020-0539-8
- InSight Marsquake Service, 2020a.** Mars Seismic Catalogue, InSight Mission; V1 2/1/2020. ETHZ, IPGP, JPL, ICL, ISAE-Supaero, MPS, Univ. Bristol. <https://doi.org/10.12686/a6>
- InSight Marsquake Service, 2020b.** Mars Seismic Catalogue, InSight Mission; V2 2020-04-01. ETHZ, IPGP, JPL, ICL, ISAE-Supaero, MPS, Univ. Bristol. <https://doi.org/10.12686/a7>
- InSight Marsquake Service, 2020c.** Mars Seismic Catalogue, InSight Mission; V3 2020-07-01. ETHZ, IPGP, JPL, ICL, ISAE-Supaero, MPS, Univ. Bristol. <https://doi.org/10.12686/a8>
- InSight Marsquake Service, 2020d.** Mars Seismic Catalogue, InSight Mission; V4 2020-10-01. ETHZ, IPGP, JPL, ICL, MPS, Univ. Bristol. <https://doi.org/10.12686/a9>
- InSight Marsquake Service, 2021a.** Mars Seismic Catalogue, InSight Mission; V5 2020-01-04. ETHZ, IPGP, JPL, ICL, MPS, Univ. Bristol. <https://doi.org/10.12686/a10>
- InSight Marsquake Service, 2021b.** Mars Seismic Catalogue, InSight Mission; V6 2021-04-01. ETHZ, IPGP, JPL, ICL, MPS, Univ. Bristol. <https://doi.org/10.12686/a11>
- InSight Marsquake Service, 2021c.** Mars Seismic Catalogue, InSight Mission; V7 2021-07-01. ETHZ, IPGP, JPL, ICL, Univ. Bristol. <https://doi.org/10.12686/a12>
- Khan, A. et al., 2021.** Upper mantle structure of Mars from InSight seismic data. Science. doi:10.1126/science.abf2966
- Kedar, S. et al., 2021.** Analyzing Low Frequency Seismic Events at Cerberus Fossae as Long Period Volcanic Quakes. <https://doi.org/10.1029/2020JE006518>
- Stähler, S. et al., 2021.** Seismic detection of the martian core. Science. doi:10.1126/science.abi7730
- van Driel, M. et al., 2021.** High frequency seismic events on Mars observed by InSight. JGR Planets. <https://doi.org/10.1029/2020JE006670>